

Thermal Stresses of Semi-Infinite Rectangular Plate: Transient Problem

Pallavi Meshram and N. W. Khobragade

Department of Mathematics, RTM Nagpur University, Nagpur (M.S), India.

Abstract- This paper is concerned with transient thermoelastic problem in which we need to determine the temperature distribution, displacement function and thermal stresses of a semi-infinite rectangular plate when the boundary conditions are known. Integral transform techniques are used to obtain the solution of the problem.

Key Words: Semi-infinite rectangular plate, transient problem, Integral transform, direct problem

I. INTRODUCTION

In 1999, **Adams and Bert** [1] studied thermoelastic vibrations of a laminated rectangular plate subjected to a thermal shock. **Tanigawa and Komatsubara** [2] discussed thermal stress analysis of a rectangular plate and its thermal stress intensity factor for compressive stress field. **Vihak; Yuzvyak and Yasinskij** [3]: derived the solution of the plane thermoelasticity problem for a rectangular domain. **Dange; Khobragade and Durge** [4] studied three dimensional inverse transient thermoelastic problem of a thin rectangular plate. **Ghume and Khobragade** [5] investigated deflection of a thick rectangular plate. **Roy and Khobragade** [6] discussed transient thermoelastic problem of an infinite rectangular slab. **Lamba and Khobragade** [7] studied thermoelastic problem of a thin rectangular plate due to partially distributed heat supply.

In 2012, **Sutar and Khobragade** [8] discussed inverse thermoelastic problem of heat conduction with internal heat generation for the rectangular plate. **Khobragade; Hiranwar; and Khalsa** [9] derived thermal deflection of a thick clamped rectangular plate. **Roy; Bagade and Khobragade** [10] studied thermal stresses of a semi infinite rectangular beam. **Jadhav and Khobragade** [11] discussed inverse thermoelastic problem of a thin finite rectangular plate due to internal heat source. **Singru and Khobragade** [12] studied thermal stress analysis of a thin rectangular plate with internal heat source. **Further Singru and Khobragade** [13] derived, Thermal stresses of a semi-infinite rectangular slab with internal heat generation.

In this paper, an attempt has been made to determine the temperature distribution, displacement function and thermal stresses functions of semi-infinite rectangular plate occupying the space $D: 0 \leq x \leq a, 0 \leq y \leq \infty$ with known boundary conditions.

II. STATEMENT OF THE PROBLEM

Consider semi-infinite rectangular plate occupying the space $D: 0 \leq x \leq a, 0 \leq y \leq \infty$. The displacement components u_x and u_y in the x and y - direction represented in the integral form as [2] are

$$u_x = \int \left[\frac{1}{E} \left(\frac{\partial^2 U}{\partial y^2} - \nu \frac{\partial^2 U}{\partial x^2} \right) + \alpha T \right] dx \quad (2.1)$$

$$u_y = \int \left[\frac{1}{E} \left(\frac{\partial^2 U}{\partial x^2} - \nu \frac{\partial^2 U}{\partial y^2} \right) + \alpha T \right] dy \quad (2.2)$$

ν and α are the Poisson's ratio and the linear coefficient of thermal expansion of the material of the plate respectively and $U(x, y, t)$ is the Airy's stress function which satisfy the following relation

$$\left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right)^2 U = -\alpha E \left(\frac{\partial^2}{\partial x^2} + \frac{\partial^2}{\partial y^2} \right) T \quad (2.3)$$

where E is the Young's modulus of elasticity and T is the temperature of the plate satisfying the differential equation

$$\frac{\partial^2 T}{\partial x^2} + \frac{\partial^2 T}{\partial y^2} = \frac{1}{k} \frac{\partial T}{\partial t} \quad (2.4)$$

subject to the initial condition

$$T(x, y, 0) = 0 \quad (2.5)$$

the boundary conditions

$$T(0, y, t) = 0 \quad (2.6)$$

$$T(a, y, t) = f(y, t) \quad (2.7)$$

$$T(x, 0, t) = 0 \quad (2.8)$$

$$T(x, \infty, t) = 0 \quad (2.9)$$

where k is the thermal diffusivity of the material of the plate.

The stress components in terms of U are given by

$$\sigma_{xx} = \frac{\partial^2 U}{\partial y^2} \tag{2.10}$$

$$\sigma_{yy} = \frac{\partial^2 U}{\partial x^2} \tag{2.11}$$

$$\sigma_{xy} = -\frac{\partial^2 U}{\partial x \partial y} \tag{2.12}$$

Equations (2.1) to (2.12) constitute the mathematical formulation of the problem under consideration.

III. SOLUTION OF THE PROBLEM

Applying Fourier sine transform to the equations (2.4), (2.5), (2.6), (2.7) and using (2.8), (2.9) one obtains

$$\frac{d^2 \bar{T}_s}{dx^2} - p^2 \bar{T}_s = \frac{1}{k} \frac{d\bar{T}_s}{dt} \tag{3.1}$$

$$\text{where } p^2 = m^2 \pi^2 \tag{3.2}$$

$$\bar{T}_s(x, m, 0) = 0 \tag{3.3}$$

$$\bar{T}_s(0, m, t) = 0 \tag{3.4}$$

$$\bar{T}_s(a, m, t) = \bar{f}_s(m, t) \tag{3.5}$$

where \bar{T}_s denotes Fourier sine transform of T and m is sine transform parameter.

Applying Laplace transform to the equations (3.1), (3.4), (3.5) and using (3.3) one obtains

$$\frac{d^2 \bar{T}_s^*}{dx^2} - q^2 \bar{T}_s^* = 0 \tag{3.6}$$

$$\text{where } q^2 = p^2 + \frac{s}{k} \tag{3.7}$$

$$\bar{T}_s^*(0, m, s) = 0 \tag{3.8}$$

$$\bar{T}_s^*(a, m, s) = \bar{f}_s^*(m, s) \tag{3.9}$$

where \bar{T}_s^* denotes Laplace transform of \bar{T}_s and s is Laplace transform parameter.

Equation (3.5) is a second order differential equation whose solution gives

$$\bar{T}_s^*(x, m, s) = Ae^{qx} + Be^{-qx} \tag{3.10}$$

where A, B are arbitrary constants.

Using (3.8) and (3.9) in (3.10) one obtains

$$A + B = 0 \tag{3.11}$$

$$Ae^{qa} + Be^{-qa} = \bar{f}_s^*(m, s) \tag{3.12}$$

Solving (3.11) and (3.12) one obtains

$$A = \frac{\bar{f}_s^*(m, s)}{e^{qa} - e^{-qa}}, \quad B = -\frac{\bar{f}_s^*(m, s)}{e^{qa} - e^{-qa}}$$

Substituting the values of A and B in (3.10) one obtains

$$\bar{T}_s^*(x, m, s) = \bar{f}_s^*(m, s) \frac{\sinh(qx)}{\sinh(qa)} \tag{3.13}$$

Applying inverse Laplace transform to the equation (3.13) one obtains

$$\bar{T}_s(x, m, t) = L^{-1} \left[\bar{f}_s^*(m, s) \frac{\sinh(qx)}{\sinh(qa)} \right] \tag{3.14}$$


To evaluate $L^{-1} \left[\bar{f}_s^*(m, s) \bar{g}_1(s) \right]$

$$\text{where } \bar{g}_1(s) = \frac{\sinh(qx)}{\sinh(qa)} \tag{3.15}$$

Using inversion integral to the equation (3.15) one obtains

$$\bar{g}_1(t) = \frac{1}{2\pi i} \int_{c-i\infty}^{c+i\infty} e^{st} \frac{\sinh(qx)}{\sinh(qa)} ds \tag{3.16}$$

Now to calculate the inversion integral (3.16) where c is greater than the real part of all singularities of the integrand. The integral is a single valued function of s in the region bounded by the closed Bromwich contour of the figure given



below:

The line NL is chosen so as to lie all the poles to the right, which are given by

$$s = s_n = \frac{in\pi}{a}, \quad n = 1, 2, \dots$$

Choosing the contour so that the curved portion LMN is an arc of the circle Γ with center at the origin and radius $R = k(n+1/2)^2(\pi/a)^2$, so that it will not pass through zero of $\sinh qa$.

The integral over the circular arc tends to zero as $n \rightarrow \infty$.

Now $\sinh(qa) = 0$ gives $q = (in\pi/a)$ ie $s = s_n = (in\pi/a)$, $n = 1, 2, 3, \dots$

Therefore

Residue at $s_n =$

$$\begin{aligned} & \lim_{s \rightarrow s_n} \left[\frac{s - s_n}{\sinh \left(a \sqrt{\frac{s}{k} + p^2} \right)} e^{st} \sinh \left(x \sqrt{\frac{s}{k} + p^2} \right) \right] \\ & = \\ & \lim_{s \rightarrow s_n} \left[\frac{2k \sqrt{\frac{s}{k} + p^2}}{a \cosh \left(a \sqrt{\frac{s}{k} + p^2} \right)} \right] \left[\lim_{s \rightarrow s_n} e^{st} \sinh \left(x \sqrt{\frac{s}{k} + p^2} \right) \right] \\ & = \frac{2kin\pi}{a^2 \cos n\pi} \left(\frac{1}{i} \right) \sin \left(ix \left(\frac{in\pi}{a} \right) \right) e^{-kt \left(p^2 + \frac{n^2\pi^2}{a^2} \right)} \end{aligned}$$

Hence the value of $\bar{g}_1(t)$ is given by

$$\bar{g}_1(t) = \frac{2k\pi}{a^2} \sum_{n=1}^{\infty} (-1)^{n+1} n \sin \left(\frac{n\pi}{a} \right) x e^{-kt \left(p^2 + \frac{n^2\pi^2}{a^2} \right)}$$

Applying the Convolution Theorem to the equation (3.14) one obtains

$$\begin{aligned} \bar{T}_s(x, m, t) &= \frac{2k\pi}{a^2} \sum_{n=1}^{\infty} (-1)^{n+1} n \sin \left(\frac{n\pi}{a} \right) x \\ & \times \int_0^t \bar{f}_s(m, t') e^{-k \left(p^2 + \frac{n^2\pi^2}{a^2} \right) (t-t')} dt' \end{aligned} \quad (3.17)$$

Applying inverse Fourier sine transform to the equations (3.17) one obtains the expressions for the temperature distribution $T(x, y, t)$ and as

$$\begin{aligned} T(x, y, t) &= \left(\frac{2k}{a^2} \right) \sum_{m=1}^{\infty} \sin py \sum_{n=1}^{\infty} (-1)^{n+1} n \sin \left(\frac{n\pi}{a} \right) x \\ & \times \int_0^t \bar{f}_s(m, t') e^{-k \left(p^2 + \frac{n^2\pi^2}{a^2} \right) (t-t')} dt' \end{aligned} \quad (3.18)$$

$$\text{where } \bar{f}_s(m, t) = \int_0^{\infty} f(y, t) \sin py dy$$

Substituting the value of $T(x, y, t)$ from (3.18) in (2.3) one obtains the expression for Airy's stress function $U(x, y, t)$ as

$$\begin{aligned} U(x, y, t) &= -\frac{\alpha E}{p^2} \left(\frac{2k}{a^2} \right) \sum_{m=1}^{\infty} \sin py \sum_{n=1}^{\infty} (-1)^{n+1} n \sin \left(\frac{n\pi}{a} \right) x \\ & \times \int_0^t \bar{f}_s(m, t') e^{-k \left(p^2 + \frac{n^2\pi^2}{a^2} \right) (t-t')} dt' \end{aligned} \quad (3.19)$$

IV. DETERMINATION OF THERMOELASTIC DISPLACEMENT

Substituting the value of $U(x, y, t)$ from (3.19) in (2.1) and (2.2) one obtains the thermoelastic displacement

functions u_x and u_y as

$$\begin{aligned} u_x &= -\left(\frac{2\alpha k}{a^3} \right) \sum_{m=1}^{\infty} \sin py \sum_{n=1}^{\infty} (-1)^{n+1} n^2 \sin \left(\frac{n\pi}{a} \right) x \\ & \times \left[2 - \frac{\nu n^2}{a^2 p^2} \right] \\ & \times \int_0^t \bar{f}_s(m, t') e^{-k \left(p^2 + \frac{n^2\pi^2}{a^2} \right) (t-t')} dt' \end{aligned} \quad (4.1)$$

$$\begin{aligned} u_y &= -\left(\frac{2\alpha k}{pa^2} \right) \sum_{m=1}^{\infty} \cos py \sum_{n=1}^{\infty} (-1)^{n+1} n \sin \left(\frac{n\pi}{a} \right) x \\ & \times \left[\frac{\pi^2 n^2}{ap^2} - \nu + 1 \right] \end{aligned}$$

$$\times \int_0^t \bar{f}_s(m, t') e^{-k \left(p^2 + \frac{n^2\pi^2}{a^2} \right) (t-t')} dt' \quad (4.2)$$

V. DETERMINATION OF STRESS FUNCTIONS

Using (3.19) in (2.10), (2.11) and (2.12) the stress functions are obtained as

$$\sigma_{xx} = \alpha E \left(\frac{2k}{a^2} \right) \sum_{m=1}^{\infty} \sin py \sum_{n=1}^{\infty} (-1)^{n+1} n \sin \left(\frac{n\pi}{a} \right) x \times \int_0^t \bar{f}_s(m, t') e^{-k \left(p^2 + \frac{n^2 \pi^2}{a^2} \right) (t-t')} dt' \quad (5.1)$$

$$\sigma_{yy} = \frac{\alpha E}{p^2} \left(\frac{2k\pi^2}{a^3} \right) \sum_{m=1}^{\infty} \sin py \sum_{n=1}^{\infty} (-1)^{n+1} n^3 \sin \left(\frac{n\pi}{a} \right) x \times \int_0^t \bar{f}_s(m, t') e^{-k \left(p^2 + \frac{n^2 \pi^2}{a^2} \right) (t-t')} dt' \quad (5.2)$$

$$\sigma_{xy} = \frac{\alpha E}{p} \left(\frac{2k\pi}{a^2} \right) \sum_{m=1}^{\infty} \cos py \sum_{n=1}^{\infty} (-1)^{n+1} n^2 \cos \left(\frac{n\pi}{a} \right) x \times \int_0^t \bar{f}_s(m, t') e^{-k \left(p^2 + \frac{n^2 \pi^2}{a^2} \right) (t-t')} dt' \quad (5.3)$$

VI. SPECIAL CASE

$$\text{Set } f(y, t) = \left(\frac{(1 - e^{-t})ya}{1 + y^2} \right) \quad (6.1)$$

Applying Fourier sine transform to the equation (6.1) one obtains

$$\bar{f}_s(m, t) = \int_0^{\infty} \left(\frac{(1 - e^{-t})ya}{1 + y^2} \right) \sin py dy = \left(1 - e^{-t} \right) \left(\frac{\pi a}{2} e^{-p} \right) \quad (6.2)$$

Substituting the value of $\bar{f}_s(m, t)$ from (6.2) in the equations (3.19) one obtains

$$T(x, y, t) = \left(\frac{\pi k}{a} \right) \sum_{m=1}^{\infty} e^{-p} \sin py \sum_{n=1}^{\infty} (-1)^{n+1} n \sin \left(\frac{n\pi}{a} \right) x$$

$$\times \int_0^t \left(1 - e^{-t'} \right) e^{-k \left(p^2 + \frac{n^2 \pi^2}{a^2} \right) (t-t')} dt' \quad (6.3)$$

VII. NUMERICAL RESULT

Set $\beta = \frac{\pi k}{\xi}$, $\pi = 3.14$, $a = 1.5$ m, $t = 1$ sec. and $k = 0.86$ in the equation (6.3) to obtain

$$\frac{T(x, y, t)}{\beta} = \sum_{m=1}^{\infty} e^{-p} \sin py \sum_{n=1}^{\infty} (-1)^{n+1} n \sin(4.2n) \times \int_0^1 \left(1 - e^{-t'} \right) e^{-0.86(2.47m^2 + 4.4n^2)(1-t')} dt' \quad (7.1)$$

VIII. CONCLUSION

In both the problems, the temperature distribution, displacement function and thermal stresses of semi-infinite rectangular plate have been investigated with the aid of integral transform techniques. The expressions are obtained in terms of Bessel's function in the form of infinite series. The results that are obtained can be applied to the design of useful structures or machines in engineering applications.

Any particular case of special interest can be derived by assigning suitable values to the parameters and functions in the expressions

REFERENCES

- [1]. **R. J. Adams and C. W Bert**, Thermoelastic vibrations of a laminated rectangular plate subjected to a thermal shock. *Journal of Thermal Stresses*, vol.22, pp. 875-895, **1999**
- [2]. **Y. Tanigawa and Y. Komatsubara**, Thermal stress analysis of a rectangular plate and its thermal stress intensity factor for compressive stress field. *Journal of Thermal Stresses*, vol.20, pp. 517-542. **1997**
- [3]. **V. M. Vihak; M. Y. Yuzvyak and A. V. Yasinskij.**: The solution of the plane thermoelasticity problem for a rectangular domain. *Journal of Thermal Stresses*, vol.21, pp. 545-561, **1998**
- [4]. **W. K. Dange; N.W. Khobragade and M. H. Durge**, Three dimensional inverse transient thermoelastic problem of a thin rectangular plate. *Int. J. of Appl. Maths*, Vol.23, No.2, 207-222, **2010**.
- [5]. **Ranjana S Ghume and N. W. Khobragade**, Deflection of a thick rectangular plate, *Canadian Journal on Science and Engg. Mathematics Research*, Vol.3 No.2, pp. 61-64, **2012**.
- [6]. **Himanshu Roy and N.W. Khobragade**, Transient thermoelastic problem of an infinite rectangular slab, *Int. Journal of Latest Trends in Maths*, Vol. 2, No. 1, pp. 37-43, **2012**.
- [7]. **N. K. Lamba; and N.W. Khobragade**, Thermoelastic problem of a thin rectangular plate due to partially distributed heat supply, *IJAMM*, Vol. 8, No. 5, pp.1-11, **2012**.

- [8]. **C. S. Sutar and N.W. Khobragade**, An inverse thermoelastic problem of heat conduction with internal heat generation for the rectangular plate, Canadian Journal of Science & Engineering Mathematics, Vol. 3, No.5, pp. 198-201, **2012**.
- [9]. **N. W. Khobragade; Payal Hiranwar; H. S. Roy and Lalsingh Khalsa**, Thermal deflection of a thick clamped rectangular plate, Int. J. of Engg. And Innovative Technology, vol. 3, Issue 1, pp. 346-348, **2013**.
- [10]. **H. S. Roy; S. H. Bagade and N. W. Khobragade**, Thermal stresses of a semi infinite rectangular beam, Int. J. of Engg. And Innovative Technology, vol. 3, Issue 1, pp. 442-445, **2013**.
- [11]. **C.M. Jadhav and N.W. Khobragade**, An inverse thermoelastic problem of a thin finite rectangular plate due to internal heat source, Int. J. of Engg. Research and Technology, vol.2, Issue 6, pp. 1009-1019, **2013**.
- [12]. **S. S. Singru and N. W. Khobragade**, Thermal stress analysis of a thin rectangular plate with internal heat source, International Journal of Latest Technology in Engineering, Management & Applied Science, Volume VI, Issue III, March 2017, 31-33.
- [13]. **S. S. Singru and N. W. Khobragade**, Thermal stresses of a semi-infinite rectangular slab with internal heat generation, International Journal of Latest Technology in Engineering, Management & Applied Science, Volume VI, Issue III, March 2017, 26-28.